

M.D. STETSON



STETCO TILE & BOWL CLEANER

Stetco Tile and Bowl Cleaner is a highly effective mild acid cleaner formulated for use in restrooms and shower rooms. Safe for use on all bathroom fixtures except wood. Easily removes rust stains, heavy soap film, lime and algae deposits. Tile and Bowl leaves any surface clean and sparkling. Use on porcelain, vitreous enamel, shower room fixtures, wash basins, toilet bowls, urinals, metal fixtures, drinking fountains, glass, copper, brass, chrome, nickel and stainless steel. Tile and Bowl is a non-fuming, non-corrosive liquid cleaner and deodorizer that leaves a pleasant scent after use.

DIRECTIONS FOR USE:

WARNING: Do not mix with bleach, ammonia or other cleaning products.

GENERAL CLEANING:

1. Apply undiluted to soiled surface with moistened mop, sponge, cloth or nozzle tip.
2. Scour with nylon abrasant pad.
3. Rinse with clean water using sponge or hose.

TILE WALLS:

1. Apply undiluted to tile surface.
2. Leave solution in place for 1 - 5 minutes.
3. Scour grout lines with nylon abrasive pad.
4. Rinse with clean water.

TOILET BOWLS & URINALS:

1. Apply undiluted to insides of bowl above water line.
2. Clean side and bottom with bowl brush or mop.
3. Flush until clean.

Please refer to Stetson's "Catch the Training" Restroom Care training guide for detailed instructions.

PACKAGING:

Available in quarts, 1 gallon jugs, 5 gallon plastic pails, 30 gallon and 55 gallon drums.

SPECIFICATIONS:

Specific Gravity.....	1.038
Color	Blue
Odor	Floral
Percent volatile by volume	83.8
pH.....	2.0 ± 0.5
Pounds/gallon	8.7 ± .2

DANGER:

DO NOT MIX WITH AMMONIA OR BLEACH. KEEP OUT OF REACH OF CHILDREN. Do not use in trigger spray bottle. Do not reuse container. Harmful if swallowed.

FIRST AID:

Avoid contact with skin and eyes. May irritate skin. Persons with cuts, allergic or sensitive skin conditions should wear rubber gloves. If contact with skin, rinse with large amounts of water. In case of contact with eyes, flush immediately with large amounts of water. Call a physician. If taken internally: Drink large amounts of water or milk of magnesia immediately. Induce vomiting. Call a physician immediately.

Refer to the material safety data sheet (MSDS) for detailed safety, health and environmental information.

March 2013